

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Ulnar Collateral Ligament Repair and Bridge Post-Operative Protocol

***AVOID ALL VALGUS STRESS**

Phase I: (0-14 days)

Immobilization in splint at 70° - MD directed

MD visit 7-14 days: Removal of splint/sutures

- Place in fully opened brace (brace for 5 weeks)

3 times per day: out of brace to work on elbow extension

Light gripping exercises

Wrist ROM

Modalities PRN

Phase II: (15-28 days)

Brace at all times except for ROM with therapist or showering

May come out of brace for ROM program

Begin elbow ROM

- A/AA ROM flexion/extension
- NO forced flexion
- Forearm pronation/supination

Begin pain free isometrics

- Wrist
- Elbow
- Shoulder (except IR/ER)

Begin manual scapular stabilization exercises

Evaluate/improve kinetic chain deficits

- Core activation
- Lower extremity flexibility/activation
- Lower body blood flow restriction training (MD approval needed)

Modalities PRN

Phase III: (29-56 days)

May come out of brace for ROM program

- Discontinue brace at 6 weeks post-op (42 days)
- If full ROM is not achieved at 6 weeks, contact physician's office

Increase ROM gradually

NO forced flexion

Begin wrist and elbow isotonic (gradual progression)

Advance scapular stabilization exercises

Advance to progressive shoulder strengthening

- Progressive ER/IR: Walkouts → ER to neutral → Full ER

Continue improving kinetic chain deficits



Ryan S. Charette MD
 Adam Q. Ferguson DO
 Jon Dale PA-C
 Christa Beyer PA-C

Aaron S. Covey MD MBA
 Rania R. Rifaey MD
 Edward S. Gensicki DPM
 Megan R. Wolf MD

Jon C. Driscoll MD
 Hudson H. Seidel MD
 Robert C. Stockton DO



www.comcllc.com
 PH: 203.265.3280
 FX: 203.741.6569

| COMPREHENSIVE PHYSICAL THERAPY | | |
|--------------------------------|---------------------------|---------------------|
| Melinda Amato OTR/L CHT | Raymond Ryan PT/Director | Peter Ives PTA |
| Brandon Heyda OTR/CHT | Valerie Peckingham PT | Kirsten Perillo PTA |
| | Isabella Mesturini PT,DPT | |

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Ulnar Collateral Ligament Repair and Bridge Post-Operative Protocol

Phase IV: (57-70 days)

Progress wrist, elbow and shoulder isotonic

- Progressive eccentric work
- Manual resistance exercises
- Advance rotator cuff strengthening to overhead position

May initiate two-handed plyometrics at 8 weeks

Phase V: (71-84 days)

Advance elbow and wrist strengthening (stress eccentrics)

Continue to progress shoulder into advanced strengthening exercises

May initiate one-handed plyometrics at 10 weeks

Phase VI: (85-112 days)

Continue to emphasize shoulder strength (rotator cuff and scapular stabilizers)

Continue endurance training

Continue kinetic chain progression

May initiate throwing mechanics drills

MD visit for possible initiation of interval throwing program

- Based on patient progression; may start around 14-18 weeks

Phase VII:

Initiate interval throwing program – MD directed

Recommend MD consult at end of flat ground throwing program

Recommend MD consult at end of mound progression

Revised 2018