

ULNAR COLLATERAL LIGAMENT REPAIR AND BRIDGE
Post-Operative Protocol

***AVOID ALL VALGUS STRESS**

Phase I: (0-14 days)

Immobilization in splint at 70°- MD directed
MD visit 7-14 days: Removal of splint/sutures

- Place in fully opened brace (brace for 5 weeks)

3 times per day: out of brace to work on elbow extension

Light gripping exercises

Wrist ROM

Modalities PRN

Phase II: (15-28 days)

Brace at all times except for ROM with therapist or showering

May come out of brace for ROM program

Begin elbow ROM

- A/AA ROM flexion/extension
- NO forced flexion
- Forearm pronation/supination

Begin pain free isometrics

- Wrist
- Elbow
- Shoulder (except IR/ER)

Begin manual scapular stabilization exercises

Evaluate/improve kinetic chain deficits

- Core activation
- Lower extremity flexibility/activation
- Lower body blood flow restriction training (MD approval needed)

Modalities PRN

Phase III: (29-56 days)

May come out of brace for ROM program

- Discontinue brace at 6 weeks post-op (42 days)
- If full ROM is not achieved at 6 weeks, contact physician's office

Increase ROM gradually

NO forced flexion

Begin wrist and elbow isotonic (gradual progression)

Advance scapular s stabilization exercises

Advance to progressive shoulder strengthening

- Progressive ER/IR: Walkouts→ER to neutral→Full ER

Continue improving kinetic chain deficits

www.meganwolfmd.com

Orthopaedics and Sports Medicine-Stratford
1901 Mooney St.
Winston-Salem, NC 27103
Phone: 336-716-8091
Fax: 336-903-7841

Orthopaedics – West Park
1917 West Park Dr.
N. Wilkesboro, NC 28659
Phone: 336-903-7845
Fax: 336-713-3244

Medical Plaza Miller
131 Miller St.
Winston-Salem, NC 27103
Phone: 336-716-8200
Fax: 336-716-9841

Phase IV: (57-70 days)

Progress wrist, elbow and shoulder isotonic

- Progressive eccentric work
- Manual resistance exercises
- Advance rotator cuff strengthening to overhead position

May initiate two-handed plyometrics at 8 weeks

Phase V: (71-84 days)

Advance elbow and wrist strengthening (stress eccentrics)

Continue to progress shoulder into advanced strengthening exercises

May initiate one-handed plyometrics at 10 weeks

Phase VI: (85-112 days)

Continue to emphasize shoulder strength (rotator cuff and scapular stabilizers)

Continue endurance training

Continue kinetic chain progression

May initiate throwing mechanics drills

MD visit for possible initiation of interval throwing program

- Based on patient progression; may start around 14-18 weeks

Phase VII:

Initiate interval throwing program – MD directed

Recommend MD consult at end of flat ground throwing program

Recommend MD consult at end of mound progression

Revised 2018