



Ryan S. Charette MD	Aaron S. Covey MD MBA	Jon C. Driscoll MD
Adam Q. Ferguson DO	Rania R. Rifaey MD	Hudson H. Seidel MD
Jon Dale PA-C	Edward S. Gensicki DPM	Robert C. Stockton DO
Christa Beyer PA-C	Megan R. Wolf MD	

A Member Of
NEOS
 NEW ENGLAND
 Orthopaedic Services
 www.comcllc.com
 PH: 203.265.3280
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY		
Melinda Amato OTR/L CHT	Raymond Ryan PT/Director	Peter Ives PTA
Brandon Heyda OTR/CHT	Valerie Peckingham PT	Kirsten Perillo PTA
	Isabella Mesturini PT,DPT	

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

1st MTP Cheilectomy Post-Operative Protocol

Weeks 0-1

- Non-weight bearing with crutch use
- Maintain and progress strength of non-operative joints (knee, hip, UE)
- Protect tissue healing: postoperative posterior splint
- Elevation above the heart “toes above the nose” for edema control

Weeks 1-4

- Weight bearing as tolerated in post-operative shoe. Wear post-op shoe with ambulation, remove at night.
- Suture removal – Week 2
- Start PT: gentle range of motion of great toe, heel raises, and toe pulls.
- Manage swelling with ice, etc.
- Do not ambulate barefoot

Weeks 4-12

- Transition to regular shoe with a wide toe box
- Continue home exercise program to include great toe range of motion
- Slowly progress low impact activity like walking
- Manage swelling with ice, etc.
- Do not ambulate barefoot

Months 3-6

- Continue progress activity as tolerated.
- Continue home exercise program
- Shoe wear should accommodate for lingering swelling.
- Do not ambulate barefoot

Revised ***