

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Operative Foot and Ankle Fracture Care

Post-operative Protocol

0-2 Weeks

- Non-weight bearing in plaster splint
- Crutches only, no scooter use
- Elevation, "toes over nose" for pain and swelling control

2-6 Weeks

- Non-weight bearing in fiberglass cast
- Okay to begin scooter use (pending physician approval)
- Elevation, "toes over nose" for pain and swelling control

6-10 Weeks

- Transition to boot, may wear a night splint during sleep
- Begin progressive weight bearing
 - Week one at 25% of weight, use two crutches
 - Week two at 50% of weight, may transition to one crutch
 - Week three may transition to full weight bearing in boot

Weeks 10-16

- Begin formal physical therapy
- At week 12, wean from boot
 - Over 10-14 days, may discontinue wearing boot at home. Continue to wear the boot outside of the home.
 - After 14 days, completely discontinue boot (14-week mark)
 - May use lace up ankle brace for activity
- May progress activity as tolerated per PT approval at 14 weeks