

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Ankle Lateral Ligament Reconstruction Post-Operative Protocol

Phase I—Maximum Protection

Weeks 0 to 2

- Post-operative posterior splint
- Non-weight bearing at all times with use of crutches
 - No scooter

Goals

- Reduce inflammation and pain
- Protect surgical repair
- Maintain strength and range of motion of non-operative joints

Exercise progression

- Open chain hip strengthening
- Gait training
- Elevation and ice to assist with swelling reduction

Phase II—Early Stretching and Strengthening

Weeks 2 to 4

- CAM boot at all times
 - Able to remove boot for ankle and foot ROM in PT
- Progress to weight bearing as tolerated
- Initiate range of motion at ankle and foot
- Initiate closed kinetic chain exercises in CAM boot
- Submaximal isometrics at ankle

Goals

- Reduce inflammation and pain
- Protect surgical repair
- Range of motion at ankle and foot
 - Avoid passive inversion and eversion
 - Avoid full-range plantarflexion (limit to 75% of ROM)
- Full weight bearing without use of crutches by 4 weeks for ambulation

Manual therapy

- Graded talocrural and subtalar mobilizations
- Passive range of motion within restrictions

Exercise progression

- Gait training
- Initiate closed chain exercises in double limb in CAM boot
- Initiate submaximal isometrics in all directions
- Elevation and ice to assist with swelling reduction

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L, CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT, DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Ankle Lateral Ligament Reconstruction Post-Operative Protocol

Phase III—Progressive Stretching and Strengthening

Weeks 4 to 8

- Transition to running shoe with Velocity ankle brace, if prescribed by MD
- Full weight bearing
- Progress to full range of motion in all directions
- Progress closed kinetic chain exercises from double limb to single limb
- Initiate balance/proprioceptive exercises

Goals

- Reduce inflammation and pain
- Protect surgical repair
- Normalized gait pattern
- Full range of motion
- Progress limb strength

Manual therapy

- Graded talocrural and subtalar mobilizations
- Passive range of motion

Exercise progression

- Normalize gait pattern
- Initiate bike and elliptical trainer
- Proprioceptive and balance drills
 - No BAPS, BOSU, or wobble board
- Unilateral closed kinetic chain strengthening program
- Modalities for pain relief and swelling reduction

Phase IV—Advance Strengthening and Running Progression

Weeks 8 to 10

- Administer Preliminary functional test at 8 weeks for physical therapist review
- Initiate straight line jogging at 8 weeks (in brace) if proper biomechanics are demonstrated and symmetry on functional test
- Initiate plyometrics in brace at 8 weeks
- Advance strengthening program

Goals

- No swelling
- Full range of motion
- Symmetrical strength and power

Exercise progression

- Basic ladder series
- Linear jogging progression
- Basic plyometric exercises starting with double limb and progressing to single limb
- Gym strengthening progression
- Progress proprioceptive exercises to varied surfaces as patient is able to demonstrate control



Ryan S. Charette MD
 Adam Q. Ferguson DO
 Jon Dale PA-C
 Christa Beyer PA-C

Aaron S. Covey MD MBA
 Rania R. Rifaey MD
 Edward S. Gensicki DPM
 Megan R. Wolf MD

Jon C. Driscoll MD
 Hudson H. Seidel MD
 Robert C. Stockton DO



www.comcllc.com
 PH: 203.265.3280
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY		
Melinda Amato OTR/L CHT	Raymond Ryan PT/Director	Peter Ives PTA
Brandon Heyda OTR/CHT	Valerie Peckingham PT	Kirsten Perillo PTA
	Isabella Mesturini PT,DPT	

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Ankle Lateral Ligament Reconstruction Post-Operative Protocol

Phase V—Return to Sport

Weeks 10 to 12

- Progress plyometric training to multi-plane, change of direction, and deceleration
- Advance strengthening program
- Administer Return to Sport functional test at 12 weeks prior to follow up appointment with MD for physician to review

Goals

- No swelling
- Full range of motion
- Normal gait pattern
- Symmetrical strength and power

Exercise progression

- Advanced ladder series
- Change of direction with running and jumping
- Sport specific field/court drills
- Gym strengthening progression

Criteria to be released for return to sport

- Follow-up examination with the physician
- Pass Return to Sport functional test at >90% (involved vs. uninvolved limb)
- Display symmetry and confidence in high-speed cutting, multi-plane plyometric drills, sprinting and decelerating

Anticipated return to sport

- 12 weeks for contact and non-contact athletes

Revised ***