

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT,DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD

### Sports Medicine/Foot & Ankle

### Osteochondral Defect of Ankle Post-Operative Protocol

#### Week 0-1

- Non-weight bearing with crutches, no scooter use
- Elevation above the heart "toes above the nose" for edema control

#### Week 1-6

##### Treatment:

- Non-weight bearing in pneumatic walking boot
- May use knee scooter
- Active ankle ROM all directions
- Begin ankle strengthening exercises
- Isometrics all directions-submaximal
- Intrinsic foot strengthening
  - toe curls, marble pickups, etc.

#### Week 6-10

##### Goals:

- Progress ankle ROM and strength
- Progressive weight bearing as follows:
  - Week 6: 25-50%
  - Week 7: 50-100%
  - Week 8: FWB
  - Weeks 8-10, wean from boot (FWB)

#### Week 10-12

##### Goals:

- Normalized gait
- Closed kinetic chain LE exercises
- Ankle ROM and strengthening
- Normal gait in shoe
- Introduce proprioceptive exercises
  - Even ground only, no BAPS, BOSU or Wobble board

##### Treatment:

- Ankle ROM
- UBE, may introduce LE stationary bike with no resistance

#### Week 12-16

##### Goals:

- Protect healing cartilage
- Return to full normal everyday activities
- Progress ankle strength, ROM and proprioceptive exercises
- Continue general strengthening of LE, UE, and trunk as indicated

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT,DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD

### Sports Medicine/Foot & Ankle

### Osteochondral Defect of Ankle Post-Operative Protocol

#### Treatment:

- Add LE closed chain exercises, single plane
- Evaluate LE biomechanics, flexibility, and strength bilaterally
- Modalities to control/decrease inflammation/pain
- Closed kinetic chain LE exercises with added weight
- Stationary bike-may start to progress resistance
- Increase duration/intensity as tolerated

#### **Week 16-20**

##### Goals:

- Patient demonstrates fair static and dynamic neuromuscular control
- Treatment:
- Introduce jogging in pool (chest deep water) or with use of Alter G
- Pre-requisites: no walking pain, adequate strength, Full ROM
- Progress proprioceptive training
- Static and dynamic balance progressing to varied surfaces as patient is able
- Cardiovascular training: continue exercise bike/elliptical

#### **Week 20-24**

##### Goals:

- Patient to demonstrate full strength throughout full ROM
- Patient to demonstrate fully restored static and dynamic neuromuscular control and kinesthetic awareness
- Treatment:
- Increase intensity of exercise bike, elliptical, stair-stepper, and treadmill
- Increase intensity and resistance in closed chain activities
- Multi-planar movements

#### **Week 24-32**

##### Goals:

- Begin high impact activity and progress as tolerated
- Running/jogging, plyometrics
- Sport specific drills
- Pass Return to sport functional test (7-8 months post-operatively)

Revised \*\*\*