

**COMPREHENSIVE PHYSICAL THERAPY**

Melinda Amato OTR/L CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT,DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD Sports Medicine/Foot & Ankle

## Hip Arthroscopy-Bilateral Labral Repair, Femoroplasty, Chondroplasty, Acetabuloplasty Post-Operative Protocol

### Phase I – Maximum Protection

#### **Weeks 0 to 3:**

- 4 point gait x 3 weeks, then wean off crutches
- Lie on stomach 2 or more hours per day
- Range of motion restrictions x 3 weeks
  - Flexion 0° to 90° for 2 weeks progressing to 120° by week 3
  - Extension 0°
  - External rotation 0°
  - Internal rotation - work for full range at 0° and 90°
  - Abduction 0° to 45°

#### Exercise progression (POD 1 to 7)

- Stationary bike with no resistance: immediately as tolerated
- Glute, quadriceps, hamstring, abduction, adduction isometrics (2x/day): immediately as tolerated
- Hip PROM (2x/day) flexion, abduction, IR supine at 90° and prone at 0°
- Hip circumduction

#### Exercise progression (POD 8 to 14)

- Hip IR/ER isometrics (2x/day)
- Initiate basic core: pelvic tilting, TVA and breathing re-education
- Beginning POD 14: quadruped rocking

#### Exercise progression (POD 15 to 21)

- Standing abduction/adduction - full weightbearing on uninvolved side only

#### **Criteria for progression to Phase 2:**

- Mobility within limitations
- Early restoration of neuromuscular control
- Normal patellar mobility

### Phase II – Progressive Stretching and Early Strengthening

#### **Weeks 3 to 6:**

- May begin deep water pool walking at 3 weeks if incisions closed, flutter/dolphin kick at 6 weeks

#### Goals

- Wean off crutches (over 7 to 10 days)
- Normal gait
- Normal single limb stance
- Full range of motion
- Improve lower extremity muscle activation, strength and endurance

#### Manual therapy

- Scar mobilization
- STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators
- Continue work on range of motion (FABER, flexion, abduction, IR, ER)

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L, CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT, DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD Sports Medicine/Foot & Ankle

## Hip Arthroscopy-Bilateral Labral Repair, Femoroplasty, Chondroplasty, Acetabuloplasty Post-Operative Protocol

### Exercise progression (as tolerated)

- Bridging double and single
- Supine dead bug series
- Sidelying hip abduction
- Quadruped hip extension series
- Standing open and closed chain multi-plane hip
- Standing internal/external rotation strengthening (use stool)
- Step-up progression
- Squat progression
- Heel raises
- Stationary biking
- Stretching: quadriceps, piriformis and hamstrings

### Criteria for progression to Phase 3:

- Hip abduction strength 4/5
- Flexion, ER and IR range of motion within normal limits
- 50% FABER range of motion compared to uninvolved side
- Normal gait
- No Trendelenberg with single leg stance/descending stairs
- Normal bilateral squat

### Phase III – Advanced Strengthening and Endurance Training

#### Weeks 6 to 12:

**Please do not discharge patient prior to 3 months without approval from Dr. Wolf**

#### Manual therapy

- STM as needed - particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking ER or FABER range of motion
- May begin trigger point dry needling for glutes, quads, adductors
  - No hip flexor tendon until week 8.
- Assess FMA and begin to address movement dysfunctions

#### Exercise progression

- Continue with muscle activation series (quadruped or straight leg series)
- Introduce movement series to increase proprioception, balance, and functional flexibility
- Progress core program as appropriate
- Advanced glute and posterior chain strengthening
- Leg press and leg curl
- Squat progression (double to single leg - add load as tolerated)
- Lunge progression
- Step-up progression
- Walking program
- Week 6:
  - Outdoor biking
  - Pool running program (at least 75% unloaded)
- Week 8 (if range of motion adequate): swimming - breast stroke kick



Ryan S. Charette MD  
 Adam Q. Ferguson DO  
 Jon Dale PA-C  
 Christa Beyer PA-C

Aaron S. Covey MD MBA  
 Rania R. Rifaey MD  
 Edward S. Gensicki DPM  
 Megan R. Wolf MD

Jon C. Driscoll MD  
 Hudson H. Seidel MD  
 Robert C. Stockton DO



www.comcllc.com  
 PH: 203.265.3280  
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY		
Melinda Amato OTR/L CHT	Raymond Ryan PT/Director	Peter Ives PTA
Brandon Heyda OTR/CHT	Valerie Peckingham PT	Kirsten Perillo PTA
	Isabella Mesturini PT,DPT	

## Megan R. Wolf, MD Sports Medicine/Foot & Ankle

## Hip Arthroscopy-Bilateral Labral Repair, Femoroplasty, Chondroplasty, Acetabuloplasty Post-Operative Protocol

### Criteria for progression to Phase 4:

- 12 weeks post-op
- Hip abduction and extension strength 5/5
- Single leg squat symmetrical with uninvolved side
- Full range of motion
- No impingement with range of motion

### Phase IV – Return to Sport Program

#### **Weeks 12 to 20:**

- May begin elliptical and stair climber at 12 weeks
- May begin return to run program if phase 4 criteria are met

#### Manual therapy

- STM as needed particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking ER or FABER range of motion
- Trigger point dry needling for glutes, TFL, quads, adductors, iliopsoas, iliacus
  - May continue to benefit patients with tightness or mild range of motion restrictions

#### Exercise progression

- Maintain muscle activation series, trunk, hip and lower extremity strength and flexibility program
- Introduce and progress plyometric program
- Begin ladder drills and multidirectional movement
- Begin interval running program
- Field/court sports specific drills in controlled environment
- Pass sports test
- Non-contact drills and scrimmaging – must have passed sports test - refer to specific return to sport program
- Return to full activity – per physician and passing PT sport test

Revised 8/2016