

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Core Muscle Injury Repair-6 Week Post-Operative Protocol

Day 1 to 7:

- Control post-operative pain and swelling. Compression shorts may be beneficial
- Avoid activity that increases intra-abdominal pressure – such as lifting.
- Recommend wearing compression shorts to help control swelling
- Walk as tolerated – aim for 1+ mile per day, increase as tolerated

Exercises:

- Kegel exercises 2 times per day
- Isometric hip adduction, glute sets, transverse abdominus (TA) activation
- Heel raises
- Light core exercises – quadruped progression, double legged bridging, clamshells

Day 8 to 14:

- Recommend wearing compression shorts to help control swelling as needed in daily activities, for exercise
- Balance core and back, anterior and posterior chain exercises
- Avoid upper body lifting
- Begin gentle soft tissue massage of abdominal muscles and hip adductors both in rehab and at home at least twice per day. May also address surrounding hip musculature.
 - Avoid aggressive massage techniques such as instrument assisted or active release

Exercises:

- If a pool is available and incision is well healed, may begin pool walking
- Progress to AROM and then light resistance exercises. Modify exercises as needed for pain, focus on good form and core control.
 - Progress clamshell and bridging resistance
 - Curl ups with TA activation. No full sit-ups.
 - Side-lying hip abduction and adduction. Start with AAROM if needed.
 - Single leg balance progression
 - Lunge progression with good form
 - Monster walks with light resistance band
 - Modified prone and side planks. Must be able to do 5 reps for 30 seconds with proper form before progressing to full planks.
 - Scapular stabilization exercises in standing
- Light stretching
 - Hip flexors
 - Hamstrings
 - Quads
 - Groin
 - Side bending
- Cardiovascular
 - Progress walking program
 - Elliptical or stair master as tolerated
 - If sufficient strength, initiate walk to jog program at day 10-12.

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L, CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT, DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Core Muscle Injury Repair-6 Week Post-Operative Protocol

Day 15-21:

- Recommend wearing compression shorts for exercise
- Balance core and back, anterior and posterior chain exercises
- Maintain proper pelvic and lower extremity alignment and control
- Continue massage of abdominals, hip adductors, as well as other hip and core musculature.
 - Avoid active release or instrument assisted massage to the hip adductors, rectus abdominus
 - Foam rollers should not be used for at least 6 weeks on hip adductors. Foam rolling of surrounding hip muscles can begin at 4 weeks post-op.

Exercises:

- Pool exercises – walking, hip AROM, swimming (no breast stroke)
- Continue single leg balance progression
- Lunge progression with good form
- Step-ups and step downs
- Minisquats progressing to full squats
- Monster walks
- Modified prone and side planks. Must be able to do 5 reps for 30 seconds with proper form before progressing to full planks.
- Continue to progress jogging program
- Initiate low impact agility program, progress as tolerated
- Initiate light sport specific activity

Day 22-28:

- Balance core and back, anterior and posterior chain exercises
- Maintain proper pelvic and lower extremity alignment and control
- Re-introduce weight training. Start at no more than 50% of pre-surgery weight level. Upper body exercises should start at/below shoulder height and then progress to overhead.
- Continue massage of abdominals, hip adductors, as well as other hip and core musculature.
 - Avoid active release or instrument assisted massage to the hip adductors, rectus abdominus
 - Foam rollers should not be used for at least 6 weeks on hip adductors. Foam rolling of surrounding hip muscles can begin at 4 weeks post-op.

Exercises:

- Pool exercises – running, agilities
- Lunge progression with good form – add multiplanar, add weight
- Single leg dead lift
- Progress agility work. Increase speed as form improves.
- Progress sport specific work into simulated non-contact drills
- Start abdominal exercises. Limit crunches and situps.



Ryan S. Charette MD
 Adam Q. Ferguson DO
 Jon Dale PA-C
 Christa Beyer PA-C

Aaron S. Covey MD MBA
 Rania R. Rifaey MD
 Edward S. Gensicki DPM
 Megan R. Wolf MD

Jon C. Driscoll MD
 Hudson H. Seidel MD
 Robert C. Stockton DO



www.comcllc.com
 PH: 203.265.3280
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY		
Melinda Amato OTR/L CHT	Raymond Ryan PT/Director	Peter Ives PTA
Brandon Heyda OTR/CHT	Valerie Peckingham PT	Kirsten Perillo PTA
	Isabella Mesturini PT,DPT	

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Core Muscle Injury Repair-6 Week Post-Operative Protocol

Day 29-35:

- Balance core and back, anterior and posterior chain exercises
- Maintain proper pelvic and lower extremity alignment and control

- Continue massage of abdominals, hip adductors, as well as other hip and core musculature as needed.
 - Avoid active release or instrument assisted massage to the hip adductors, rectus abdominus
 - Foam rollers should not be used for at least 6 weeks on hip adductors. Foam rolling of surrounding hip muscles can begin at 4 weeks post-op.

Exercises:

- Single leg hops
- Box Jumps
- Continue sport specific exercises
- Initiate participation in individual drills in practice, progress to full participation as tolerated

Day 36+:

- Progress into full sport activity
- Continue long term core strengthening program
 - Avoid active release or instrument assisted massage to the hip adductors, rectus abdominus

Revised ***