

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L, CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT, DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Endoscopic Iliotibial Band Recession Post-Operative Protocol

Progress as tolerated through the following phases

Phase I – Maximum Protection

Weeks 1 to 2:

- 50% weightbearing for 2 weeks
- No brace
- Range of motion restrictions for 3 weeks: no end range of motion stretching in flexion or extension

Exercise progression (POD 1 to 7)

- Stationary bike with no resistance: immediately as tolerated
- Glute, quadriceps, hamstring, abduction, adduction isometrics (2x/day): immediately as tolerated
- Hip circumduction

Exercise progression (POD 8 to 14)

- Hip IR/ER isometrics (2x/day)
- Initiate basic core: pelvic tilting, TVA and breathing re-education
- Quadruped rocking

Exercise progression (POD 15 to 21)

- Standing abduction/adduction - full weightbearing on uninvolved side only

Criteria for progression to Phase 2:

- Mobility within limitations
- Early restoration of neuromuscular control
- Normal patellar mobility

Phase II – Progressive Stretching and Early Strengthening

- May begin deep water pool walking at 3 weeks if incisions closed.

Goals

- Wean off crutches (over 7 to 10 days)
- Normal gait
- Normal single limb stance
- Full range of motion
- Improve lower extremity muscle activation, strength and endurance

Manual therapy

- Scar mobilization
- STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators
- Continue work on range of motion (FABER, flexion, abduction, IR, ER)

Exercise progression (as tolerated)

- Bridging double and single
- Supine dead bug series
- Sidelying hip abduction
- Quadruped hip extension series
- Standing internal/external rotation strengthening (use stool)
- Step-up progression
- Squat progression
- Heel raises
- Stationary biking
- Stretching: quadriceps, piriformis and hamstrings

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Endoscopic Iliotibial Band Recession Post-Operative Protocol

Criteria for Progression to Phase 3:

- Hip abduction strength 4/5
- Flexion, ER and IR range of motion within normal limits
- Normal gait
- No Trendelenberg with single leg stance/descending stairs
- Normal bilateral squat

Phase III – Advanced Strengthening and Endurance Training

Manual therapy

- Continue soft tissue mobilization as needed particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking ER or FABER range of motion
- May begin trigger point dry needling for glutes, quads, adductors.
- Assess FMA and begin to address movement dysfunctions

Exercise progression

- Continue with muscle activation series (quadruped or straight leg series)
- Introduce movement series to increase proprioception, balance, and functional flexibility
- Progress core program as appropriate
- Standing open and closed chain multi-plane hip
- Advanced glute and posterior chain strengthening
- Leg press and leg curl
- Squat progression (double to single leg - add load as tolerated)
- Lunge progression
- Step-up progression
- Walking program
- Outdoor biking
- Swimming
- Pool running program (at least 75% unloaded)

Criteria for progression to Phase 4:

- Hip abduction and extension strength 5/5
- Single leg squat symmetrical with uninvolved side
- Full range of motion
- No impingement with range of motion

Phase IV – Return to Sport Program

- May begin return to run program if phase 4 criteria are met

Manual Therapy

- Continue soft tissue mobilization as needed particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking end range FABER range of motion
- Trigger point dry needling for glutes, TFL, quads, adductors, iliopsoas, iliacus
 - May continue to benefit patients with tightness or mild range of motion restrictions



Ryan S. Charette MD
Adam Q. Ferguson DO
Jon Dale PA-C
Christa Beyer PA-C

Aaron S. Covey MD MBA
Rania R. Rifaey MD
Edward S. Gensicki DPM
Megan R. Wolf MD

Jon C. Driscoll MD
Hudson H. Seidel MD
Robert C. Stockton DO



www.comcllc.com
PH: 203.265.3280
FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Endoscopic Iliotibial Band Recession Post-Operative Protocol

Exercise progression

- Maintain muscle activation series, trunk, hip and lower extremity strength and flexibility program
- Introduce and progress plyometric program
- Begin ladder drills and multidirectional movement
- Begin interval running program
- Field/court sports specific drills in controlled environment
- Pass sports test
- Non-contact drills and scrimmaging – must have passed sports test - refer to specific return to sport program
- Return to full activity – per physician and passing PT sport test

Revised ***