

**COMPREHENSIVE PHYSICAL THERAPY**

Melinda Amato OTR/L, CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT, DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD

### Sports Medicine/Foot & Ankle

### Hip Arthroscopy with Core Decompression or Microfracture Post-Operative Protocol

#### Phase I – Maximum Protection

##### Weeks 0 to 6:

- Toe-touch weightbearing for 6 weeks
- Lie on stomach 2 or more hours per day
- CPM 4-6 hours per day x 6 weeks

##### Manual therapy

- Scar mobilization once incisions are closed
- STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators

##### Exercise progression (POD 1 to 7)

- Stationary bike with no resistance: immediately as tolerated
- Glute, quadriceps, hamstring isometrics, abduction, adduction (2x per day): immediately as tolerated

##### Exercise progression (POD 8 to 21)

- Hip isometrics IR/ER (2x/day)
- Initiate basic core: pelvic tilting, TVA and breathing re-education

##### Exercise progression (POD 21 to Week 6)

- Non-weightbearing strength during this phase
- Standing abduction/adduction/extension/flexion - full weightbearing on uninvolved side only
- Supine dead bug series
- Introduce basic core program
- Sidelying hip abduction
- Quadruped hip extension series
- Week 5: may begin quadruped rocking for flexion range of motion

##### **Criteria for progression to Phase 2:**

- Mobility within limitations
- Early restoration of neuromuscular control and non-weight bearing strengthening
- Normal patellar mobility

#### Phase II – Progressive Stretching and Early Strengthening

##### Weeks 6 to 9:

##### Goals

- Wean off crutches (over 7 to 10 days)
- Normal gait
- Normal single limb stance
- Full range of motion
- Improve lower extremity muscle activation, strength and endurance

##### Manual therapy

- Scar mobilization
- STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators
- Continue work on range of motion (FABER, flexion, abduction, IR, ER)

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT,DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD

### Sports Medicine/Foot & Ankle

### Hip Arthroscopy with Core Decompression or Microfracture Post-Operative Protocol

#### Exercise progression (as tolerated)

- Bridging double and single
- Supine dead bug series
- Sidelying hip abduction
- Quadruped hip extension series
- Standing open and closed chain multi-plane hip
- Standing internal/external rotation strengthening (use stool)
- Step-up progression
- Squat progression
- Heel raises
- Stationary biking
- Deep water pool program when incisions are completely healed
- Stretching: quadriceps, piriformis and hamstrings

#### **Criteria for Progression to Phase 3:**

- Hip abduction strength 4/5
- Flexion, ER and IR range of motion within normal limits
- 50% FABER range of motion compared to uninvolved side
- Normal gait
- No Trendelenberg with single leg stance/descending stairs
- Normal bilateral squat

#### **Phase III – Advanced Strengthening and Endurance Training**

##### **Weeks 9 to 16:**

**Please do not discharge patient prior to 3 months without approval from Dr. Wolf**

#### Manual therapy

- STM as needed - particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking ER or FABER range of motion
- May begin trigger point dry needling for glutes, quads, adductors
- Assess FMA and begin to address movement dysfunctions

#### Exercise progression

- Continue with muscle activation series (quadruped or straight leg series)
- Introduce movement series to increase proprioception, balance, and functional flexibility
- Progress core program as appropriate
- Advanced glute and posterior chain strengthening
- Leg press and leg curl
- Squat progression (double to single leg - add load as tolerated)
- Lunge progression
- Step-up progression
- Walking program
- Week 9
  - Outdoor biking
  - Swimming: breast stroke kick
  - Shallow water pool running program - 75% unloaded



Ryan S. Charette MD  
 Adam Q. Ferguson DO  
 Jon Dale PA-C  
 Christa Beyer PA-C

Aaron S. Covey MD MBA  
 Rania R. Rifaey MD  
 Edward S. Gensicki DPM  
 Megan R. Wolf MD

Jon C. Driscoll MD  
 Hudson H. Seidel MD  
 Robert C. Stockton DO



www.comcllc.com  
 PH: 203.265.3280  
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY		
Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT	Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT	Peter Ives PTA Kirsten Perillo PTA

## Megan R. Wolf, MD Sports Medicine/Foot & Ankle

## Hip Arthroscopy with Core Decompression or Microfracture Post-Operative Protocol

### Criteria for progression to Phase 4:

- 12 weeks post-op
- Hip abduction and extension strength 5/5
- Single leg squat symmetrical with uninvolved side
- Full range of motion
- No impingement with range of motion

### Phase IV – Return to Sport Program

#### **Weeks 16 to 24:**

- May begin elliptical and stair climber at 16 weeks
- May begin return to run program if phase 4 criteria are met

#### Exercise progression

- Maintain muscle activation series, trunk, hip and lower extremity strength and flexibility program
- Introduce and progress plyometric program
- Begin ladder drills and multidirectional movement
- Begin interval running program
- Field/court sports specific drills in controlled environment
- Pass sports test
- Non-contact drills and scrimmaging – must have passed sports test - refer to specific return to sport program
- Return to full activity – per physician and therapist

Revised 8/2016