

**COMPREHENSIVE PHYSICAL THERAPY**

Melinda Amato OTR/L, CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT, DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD

### Sports Medicine/Foot & Ankle

### Hip Arthroscopy-Labral Debridement Post-Operative Protocol

Progress as tolerated through the following phases

#### Phase I – Maximum Protection

##### **Weeks 1 to 2:**

- 50% weightbearing for 2 weeks
- Range of motion restrictions for 3 weeks: no end range of motion stretching in flexion or extension

##### Exercise progression (POD 1 to 7)

- Stationary bike with no resistance: immediately as tolerated
- Glute, quadriceps, hamstring, abduction, adduction isometrics (2x/day): immediately as tolerated
- Hip circumduction

##### Exercise progression (POD 8 to 14)

- Hip IR/ER isometrics (2x/day)
- Initiate basic core: pelvic tilting, TVA and breathing re-education
- Quadruped rocking

##### Exercise progression (POD 15 to 21)

- Standing abduction/adduction - full weightbearing on uninvolved side only

#### **Criteria for progression to Phase 2:**

- Mobility within limitations
- Early restoration of neuromuscular control
- Normal patellar mobility

#### Phase II – Progressive Stretching and Early Strengthening

- May begin deep water pool walking at 3 weeks if incisions closed.

##### Goals

- Wean off crutches (over 7 to 10 days)
- Normal gait
- Normal single limb stance
- Full range of motion
- Improve lower extremity muscle activation, strength and endurance

##### Manual therapy

- Scar mobilization
- STM to quad, ITB, hip flexors, glutes, hip adductors/abductors/rotators
- Continue work on range of motion (FABER, flexion, abduction, IR, ER)

##### Exercise progression (as tolerated)

- Bridging double and single
- Supine dead bug series
- Sidelying hip abduction
- Quadruped hip extension series
- Standing internal/external rotation strengthening (use stool)
- Step-up progression
- Squat progression
- Heel raises
- Stationary biking
- Stretching: quadriceps, piriformis and hamstrings

**COMPREHENSIVE PHYSICAL THERAPY**

Melinda Amato OTR/L CHT  
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director  
Valerie Peckingham PT  
Isabella Mesturini PT,DPT

Peter Ives PTA  
Kirsten Perillo PTA

## Megan R. Wolf, MD

### Sports Medicine/Foot & Ankle

### Hip Arthroscopy-Labral Debridement Post-Operative Protocol

#### Criteria for Progression to Phase 3:

- Hip abduction strength 4/5
- Flexion, ER and IR range of motion within normal limits
- Normal gait
- No Trendelenberg with single leg stance/descending stairs
- Normal bilateral squat

#### Phase III – Advanced Strengthening and Endurance Training

##### Manual therapy

- Continue soft tissue mobilization as needed particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking ER or FABER range of motion
- May begin trigger point dry needling for glutes, quads, adductors.
- Assess FMA and begin to address movement dysfunctions

##### Exercise progression

- Continue with muscle activation series (quadruped or straight leg series)
- Introduce movement series to increase proprioception, balance, and functional flexibility
- Standing open and closed chain multi-plane hip
- Advanced glute and posterior chain strengthening
- Leg press and leg curl
- Squat progression (double to single leg - add load as tolerated)
- Lunge progression
- Step-up progression
- Walking program
- Outdoor biking
- Swimming
- Pool running program (at least 75% unloaded)

#### Criteria for progression to Phase 4:

- Hip abduction and extension strength 5/5
- Single leg squat symmetrical with uninvolved side
- Full range of motion
- No impingement with range of motion



|                     |                        |                       |
|---------------------|------------------------|-----------------------|
| Ryan S. Charette MD | Aaron S. Covey MD MBA  | Jon C. Driscoll MD    |
| Adam Q. Ferguson DO | Rania R. Rifaey MD     | Hudson H. Seidel MD   |
| Jon Dale PA-C       | Edward S. Gensicki DPM | Robert C. Stockton DO |
| Christa Beyer PA-C  | Megan R. Wolf MD       |                       |

A Member Of  
**NEOS**  
 NEW ENGLAND  
 Orthopaedic Services  
 www.comcllc.com  
 PH: 203.265.3280  
 FX: 203.741.6569

**COMPREHENSIVE PHYSICAL THERAPY**

|                         |                           |                     |
|-------------------------|---------------------------|---------------------|
| Melinda Amato OTR/L CHT | Raymond Ryan PT/Director  | Peter Ives PTA      |
| Brandon Heyda OTR/CHT   | Valerie Peckingham PT     | Kirsten Perillo PTA |
|                         | Isabella Mesturini PT,DPT |                     |

# Megan R. Wolf, MD

## Sports Medicine/Foot & Ankle

## Hip Arthroscopy-Labral Debridement Post-Operative Protocol

### Phase IV – Return to Sport Program

- May begin return to run program if phase 4 criteria are met

#### Manual Therapy

- Continue soft tissue mobilization as needed particularly glutes, adductors, hip flexors, abductors
- Gentle joint mobilizations as needed for patients lacking end range FABER range of motion
- Trigger point dry needling for glutes, TFL, quads, adductors, iliopsoas, iliacus
  - May continue to benefit patients with tightness or mild range of motion restrictions

#### Exercise progression

- Maintain muscle activation series, trunk, hip and lower extremity strength and flexibility program
- Introduce and progress plyometric program
- Begin ladder drills and multidirectional movement
- Begin interval running program
- Field/court sports specific drills in controlled environment
- Pass sports test
- Non-contact drills and scrimmaging – must have passed sports test - refer to specific return to sport program
- Return to full activity – per physician and passing PT sport test

Revised 8/2016