

Megan R. Wolf, MD Sports Medicine

ANTERIOR CRUCIATE LIGAMENT (ACL) RECONSTRUCTION WITH MENISCUS TRANSPLANT

Post-Operative Protocol

Phase I - Maximum Protection

Weeks 0 to 6:

- Brace 0-90 x 6 weeks
 - Recommend locking in extension for sleep
- Non-weight bearing with two crutches for 6 weeks
- Limit knee flexion to 90-degrees for 3 weeks; allow 90-120 degrees between 3 to 6 weeks

Goals

- o Reduce inflammation and pain
- o 0 degrees of knee extension

Exercise progression

- Quadriceps setting using NMES as needed
- Emphasize patellofemoral mobilizations
- Passive/active knee range of motion with 90° flexion limit
- Quadriceps setting emphasize VMO function
- o Multi-plane straight leg raising
- Open chain hip strengthening
- Gait training

Phase II - Progressive Stretching and Early Strengthening

Weeks 6 to 8:

- Brace open to 0-90 degrees for ambulation
- Continue using two crutches, with gradual progression of weight bearing
 - Increase WB to FWB over next 2 weeks

Exercise progression

- o Gradual progression to full knee flexion
- o Continue to emphasize patella mobility
- Begin bilateral closed kinetic chain strengthening limited range initially
- Step-up progression
- Begin stationary bike with light resistance initially
- o Gait training normalize gait pattern

Phase III - Advanced Strengthening and Endurance Training

Weeks 8 to 10:

Goals

Full knee range of motion

Exercise progression

- o Avoid rotational movements for 14 weeks
- Advance stationary biking program (increase intensity)
- Introduce treadmill walking and elliptical trainer
- Begin unilateral closed kinetic chain program
- o Gym strengthening progression (leg press above 90-degrees, hamstrings curls etc.)

Fax: 336-713-3244

Phone: 336-716-8200 Fax: 336-716-9841

Weeks 10 to 12:

Exercise progression

- Outdoor biking
- o Lunge progression (retro, walk and split) as indicated
- o Forward/backward elevated treadmill walking
- Deep water pool running progression

Weeks 12-14

Administer preliminary functional test for physician to review

<u>Phase IV – Advance Strengthening and Running Progression</u>

Weeks 12 to 20:

Exercise progression

- o Progress resistance with squat and lunge strengthening program.
- o Begin light plyometric drills, progress from bilateral to unilateral
- o Begin linear jogging week 16
- Progress to lateral and rotational stresses at 18 weeks
- Multi-directional drills at 18-20 weeks

Phase V – Return to Sport

Weeks 20 to 24:

Exercise progression

- o Advance ladder, hurdle and plyo box progressions
- Sport specific field/court drills
- Non-contact drills

Criteria to be released for return to sport

- o Follow-up examination with the physician
- o Pass return-to-sport functional test at >90% (involved vs uninvolved limb)
 - See testing protocol
- Display symmetry and confidence in high-speed cutting, multi-plane plyometric drills, sprinting and decelerating

Revised 9/2019