

Ryan S. Charette MD Adam Q Ferguson DO Jon Dale PA-C Christa Beyer PA-C Aaron S Covey MD MBA Rania R. Rifaey MD Edward S. Gensicki DPM Megan R. Wolf MD Jon C Driscoll MD Hudson H. Seidel MD Robert C. Stockton DO

Peter Ives PTA

Kirsten Perillo PTA

A Member Of

NEW ENGLAND

Orthopaedic Services

www.comcllc.com PH:203.265.3280 FX:203.741.6569

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT

sabelia Mesturini PI,DPI

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Anterior Cruciate Ligament (ACL) Reconstruction With Meniscus Repair (Radial/Root)

Post-Operative Protocol

Phase I – Maximum Protection

Weeks 0 to 6:

- Brace 0-90 x 6 weeks
 - o Recommend locking in extension for sleep
- Non-weight bearing with two crutches for 6 weeks
- Limit knee flexion to 90-degrees for 3 weeks; allow 90-120 degrees between 3 to 6 weeks
 Goals
 - o Reduce inflammation and pain
 - 0 degrees of knee extension

Exercise progression

- Quadriceps setting using NMES as needed
- Emphasize patellofemoral mobilizations
- o Passive/active knee range of motion with 90° flexion limit
- Quadriceps setting emphasize VMO function
- Multi-plane straight leg raising
- Open chain hip strengthening
- o Gait training

Phase II – Progressive Stretching and Early Strengthening

Weeks 6 to 8:

- Brace open to 0-90 degrees for ambulation
- Continue using two crutches, with gradual progression of weight bearing
 - Increase WB to FWB over next 2 weeks

Exercise progression

- Gradual progression to full knee flexion
- Continue to emphasize patella mobility
- Begin bilateral closed kinetic chain strengthening limited range initially
- Step-up progression
- Begin stationary bike with light resistance initially
- Gait training normalize gait pattern

Phase III - Advanced Strengthening and Endurance Training

Weeks 8 to 10:

Goals

o Full knee range of motion

Exercise progression

- Avoid rotational movements for 14 weeks
- Advance stationary biking program (increase intensity)
- Introduce treadmill walking and elliptical trainer
- o Begin unilateral closed kinetic chain program
- Gym strengthening progression (leg press above 90-degrees, hamstrings curls etc.)

PH: 203.265.3280



Ryan S. Charette MD Adam Q Ferguson DO Jon Dale PA-C Christa Beyer PA-C

Aaron S Covey MD MBA Rania R. Rifaey MD Edward S. Gensicki DPM Megan R. Wolf MD Jon C Driscoll MD Hudson H. Seidel MD Robert C. Stockton DO



www.comcllc.com PH:203.265.3280 FX:203.741.6569

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT Peter Ives PTA Kirsten Perillo PTA

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Anterior Cruciate Ligament (ACL) Reconstruction With Meniscus Repair (Radial/Root)

Post-Operative Protocol

Weeks 10 to 12:

Exercise progression

- o Outdoor biking
- Lunge progression (retro, walk and split) as indicated
- Forward/backward elevated treadmill walking
- Deep water pool running progression

Weeks 12-14

Administer preliminary functional test for physician to review

Phase IV - Advance Strengthening and Running Progression

Weeks 12 to 20:

Exercise progression

- o Progress resistance with squat and lunge strengthening program.
- Begin light plyometric drills, progress from bilateral to unilateral
- o Begin linear jogging week 16
- o Progress to lateral and rotational stresses at 18 weeks
- Multi-directional drills at 18-20 weeks

Phase V – Return to Sport

Weeks 20 to 24:

Exercise progression

- Advance ladder, hurdle and plyo box progressions
- Sport specific field/court drills
- Non-contact drills

Criteria to be released for return to sport

- o Follow-up examination with the physician
- Pass return-to-sport functional test at >90% (involved vs uninvolved limb)
 - See testing protocol
- Display symmetry and confidence in high-speed cutting, multi-plane plyometric drills, sprinting and decelerating

Revised 9/2019