



Ryan S. Charette MD
 Adam Q. Ferguson DO
 Jon Dale PA-C
 Christa Beyer PA-C

Aaron S. Covey MD MBA
 Rania R. Rifaey MD
 Edward S. Gensicki DPM
 Megan R. Wolf MD

Jon C. Driscoll MD
 Hudson H. Seidel MD
 Robert C. Stockton DO



www.comcllc.com
 PH: 203.265.3280
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY		
Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT	Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT	Peter Ives PTA Kirsten Perillo PTA

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

Articular Cartilage Restoration— Patellofemoral Compartment Post-Operative Protocol

Phase I - Maximum Protection

Weeks 0-6

- Brace locked 0-20 degrees
- Weight bearing progression with use of axillary crutches
 - Week 1: <20% of body weight
 - Weeks 2-6: progress from 20% to 50% of body weight
- Initiate quadriceps muscle activation
- Initiate range of motion (restrictions apply to unloaded and loaded motion)
 - Week 1: 0-20 degrees flexion
 - Weeks 2-3: 0-60 degrees flexion
 - Weeks 4-6: 0-125 degrees flexion
- CPM use 6 hours a day for 6 weeks
 - No brace use when using CPM
 - Range of motion on CPM consistent with ROM restrictions listed above.

Goals

- Reduce inflammation and pain
- Protect surgical repair
- Maintain full knee extension range of motion
- Gradually progress knee range of motion per above restrictions
- Maintain strength and motion of non-operative joints
- Quadriceps activation

Exercise progression

- Passive/active knee range of motion
- Quad sets, hamstrings sets, glute sets
- Multi-plane open kinetic chain strengthening (i.e. straight leg raises, avoid patellofemoral provocative exercises)
- Initiate bike with no resistance to facilitate ROM at 4 weeks
- Use of BFR (blood flow restriction) therapy to facilitate strengthening during weight bearing restrictions
- Patellofemoral mobilizations
- Gait training
- Elevation and cryotherapy to assist with swelling reduction

Phase II- Progressive Stretching and Early Strengthening

Weeks 6 to 12:

- Unlock brace at 6 weeks and discontinue once full weight bearing
- Weight bearing as tolerated progressing to full weight bearing
 - Progress to weight bearing as tolerated pending MD approval after imaging at 6 week follow up visit
- Full range of motion
- Initiate closed chain strengthening
- Initiate balance and proprioception exercises

Goals

- Reduce inflammation and pain
- Protect surgical repair
- Full knee range of motion
- Maintain strength of non-operative joints
- Full weight bearing by 8 weeks
- Normalizing gait pattern

863 N. Main St. Ext., Suite 200
 Wallingford, CT 06492

455 Lewis Avenue, Suite 101
 Meriden, CT 06451

98 Main Street, Suite 201
 Southington, CT 06489

714 South Main Street
 Cheshire, CT 06410

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT
Brandon Heyda OTR/CHT

Raymond Ryan PT/Director
Valerie Peckingham PT
Isabella Mesturini PT,DPT

Peter Ives PTA
Kirsten Perillo PTA

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Articular Cartilage Restoration— Patellofemoral Compartment Post-Operative Protocol

Exercise progression

- Able to gradually increase resistance on bike at 6 weeks
- Initiate elliptical at 10 weeks
- Initiate closed chain strengthening in double limb progressing to single limb
- Maintain squat depth at 90 degrees or above
- Step up progression
- Gait training
- Elevation and cryotherapy to assist with swelling reduction

Phase III- Progressive Strengthening

Weeks 12 to 24:

- Advance strengthening program
- Balance and proprioceptive exercises

Goals

- Reduce inflammation and pain
- Protect surgical repair
- Full knee range of motion
- Progress limb strength
- Normal gait pattern

Exercise progression

- Progress closed chain single and double limb strength as able
 - Avoid patellofemoral provocative exercises (lunges, leg extension)

Phase IV- Advanced Strengthening, Running Progression, Plyometric Training

Months 6 to 9:

- Administer Preliminary functional test at 6 months for MD to review
- Initiate straight line jogging at 6 months if proper biomechanics are demonstrated and symmetry on functional test
- Advance strengthening program
- Initiate plyometric training in double limb with gradual progression to single limb
- Able to return to low-impact recreational activities (walking, biking, elliptical, swimming)

Goals

- No swelling
- Full range of motion
- Normal gait pattern
- Symmetrical strength and power

Exercise progression

- Single limb closed chain strengthening
- Proprioception drills
- Basic ladder series
- Linear jogging progression
- Basic plyometric box progression
- Gym strengthening progression



Ryan S. Charette MD	Aaron S. Covey MD MBA	Jon C. Driscoll MD
Adam Q. Ferguson DO	Rania R. Rifaey MD	Hudson H. Seidel MD
Jon Dale PA-C	Edward S. Gensicki DPM	Robert C. Stockton DO
Christa Beyer PA-C	Megan R. Wolf MD	

A Member Of
NEOS
 NEW ENGLAND
 Orthopaedic Services
 www.comcllc.com
 PH: 203.265.3280
 FX: 203.741.6569

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT	Raymond Ryan PT/Director	Peter Ives PTA
Brandon Heyda OTR/CHT	Valerie Peckingham PT	Kirsten Perillo PTA
	Isabella Mesturini PT,DPT	

Megan R. Wolf, MD

Sports Medicine/Foot & Ankle

**Articular Cartilage Restoration—
 Patellofemoral Compartment
 Post-Operative Protocol**

Phase V- Return to Sport

Months 9 to 12:

- Progress plyometric training to single limb, multi-plane, change of direction, and deceleration
- Advance strengthening program
- Administer Return To Sport functional test prior to 12 month follow up appointment with MD for physician to review

Goals

- No swelling
- Full range of motion
- Normal gait pattern
- Symmetrical strength and power

Exercise progression

- Advanced ladder series
- Change of direction with running and jumping
- Sport specific field/court drills
- Gym strengthening progression

Criteria to be released for return to sport

- Follow-up examination with the physician
- Pass Return To Sport functional test at >90% (involved vs. uninvolved limb)
- Display symmetry and confidence in high-speed cutting, multi-plane plyometric drills, sprinting and decelerating

Anticipated return to sport:

12 months for contact and non-contact athletes

Revised ***