

# Megan R. Wolf, MD Sports Medicine

## MENISCECTOMY, DEBRIDEMENT, LYSIS OF ADHESIONS (LOA) WITH DEGENERATIVE JOINT DISEASE (DJD)

Post-Operative Protocol

#### Phase I - Maximum Protection (Weeks 0 to 2):

Use crutches weight 50% bearing x 1-2 weeks

#### Goals

- Reduce inflammation
- Normalize patella mobility with manual mobilizations
- Full ROM progressing as tolerated

#### Exercise progression

- Quadriceps setting using NMES as needed
- Multi-plane straight leg raising
- Gait training

#### Phase II – Progressive Stretching and Early Strengthening (Weeks 2 to 6):

#### Weeks 2 to 4 weeks:

#### Goals

- Progress off crutches
- Full knee ROM
- Normalize gait mechanics

#### Exercise progression

- Multi-plane open and closed kinetic chain hip strengthening
- Step-up progression
- Stationary biking and treadmill/outdoor walking
- Proprioception drills

#### Weeks 4 to 6 weeks:

#### Goals

- Control inflammation
- Full knee ROM
- Increase strength and muscular endurance

#### Exercise progression

- Leg press, mini squats (as/if tolerated), hamstrings curls
- Increase intensity of stationary bike program, may add elevation to treadmill walking and elliptical
- Deep water pool program when incisions are completely healed

#### Phase III/IV – Advanced Strengthening and Endurance Training (Weeks 6 to 12):

#### Weeks 6 to 12:

#### Goals

- Full knee flexion and extension with terminal stretch
- Progressive strengthening
- Increase muscular endurance

Fax: 336-713-3244

Phone: 336-716-8200 Fax: 336-716-9841

### Exercise progression

- Outdoor biking
- Swimming
- Elliptical
- Interval golf program

Revised \*\*\*