Ryan S. Charette MD Adam Q Ferguson DO Jon Dale PA-C Christa Beyer PA-C Aaron S Covey MD MBA Rania R. Rifaey MD Edward S. Gensicki DPM Megan R. Wolf MD Jon C Driscoll MD Hudson H. Seidel MD Robert C. Stockton DO



www.comcllc.com PH:203.265.3280 FX:203.741.6569

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT Peter Ives PTA Kirsten Perillo PTA

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Meniscus Repair (Radial/Root)

Post-Operative Protocol

Phase I – Maximum Protection

Weeks 0 to 6:

- Brace 0-90 degrees
 - Recommend locking brace in extension for sleep
- Non-weight bearing with two crutches for 6 weeks
- Limit knee flexion to 90-degrees for 3 weeks; allow 90-120 degrees between 3 to 6 weeks
 Goals
 - Reduce inflammation and pain
 - o 0 degrees of knee extension

Exercise progression

- Emphasize patellofemoral mobilizations
- Passive/active knee range of motion with 90° flexion limit
- Quadriceps setting emphasize VMO function
- o Multi-plane straight leg raising
- Open chain hip strengthening
- Gait training

Phase II - Progressive Stretching and Early Strengthening

Weeks 6 to 8:

- Brace open to 0-90 degrees for ambulation
- Continue using two crutches, with gradual progression of weight bearing
 - Increase WB to FWB over next 2 weeks

Exercise progression

- o Gradual progression to full knee flexion
- o Continue to emphasize patella mobility
- Begin bilateral closed kinetic chain strengthening limited range initially
- Step-up progression
- o Begin stationary bike with light resistance initially
- Gait training normalize gait pattern

Phase III - Advanced Strengthening and Proprioception

Weeks 8 to 12:

Goals

Full knee range of motion

Exercise progression

- Avoid rotational movements for 14 weeks
- Advance stationary biking program (increase intensity)
- Introduce treadmill walking and elliptical trainer
- Begin unilateral closed kinetic chain program
- o Gym strengthening progression (leg press above 90-degrees, hamstrings curls etc.)

863 N. Main St. Ext., Suite 200 Wallingford, CT 06492 455 Lewis Avenue, Suite 101 Meriden, CT 06451 98 Main Street, Suite 201 Southington, CT 06489

FX: 203.741.6569

714 South Main Street Cheshire, CT 06410



Ryan S. Charette MD Adam Q Ferguson DO Jon Dale PA-C Christa Beyer PA-C Aaron S Covey MD MBA Rania R. Rifaey MD Edward S. Gensicki DPM Megan R. Wolf MD Jon C Driscoll MD Hudson H. Seidel MD Robert C. Stockton DO



www.comcllc.com PH:203.265.3280 FX:203.741.6569

COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT

Kirsten Perillo PTA

Peter Ives PTA

Megan R. Wolf, MD Sports Medicine/Foot & Ankle

Meniscus Repair (Radial/Root)

Post-Operative Protocol

<u>Phase IV – Advanced Strengthening and Plyometric Drills</u> Weeks 12 to 16:

Exercise progression

Begin light bilateral plyometrics

Phase V - Return to Sport

Weeks 16-20:

Exercise Progression

- Plyometric drills from bilateral to unilateral at 16 weeks
- Linear running progression at 16 weeks
- o Progress to lateral and rotational stresses at 18 weeks
- Multi-directional drills at 18-20 weeks

Criteria to be released for return to sport

- o Follow-up examination with the physician
- Pass return to sport functional test at >90% (involved vs uninvolved limb)
 - Perform test at 20 weeks or after, depending on patient readiness
 - See testing protocol
- Display symmetry and confidence in high-speed cutting, multi-plane plyometric drills, sprinting and decelerating

Revised 9/2019