

**SHOULDER MANIPULATION UNDER ANESTHESIA**  
Post-Operative Protocol

---

**Phase I – Maximum Protection – Passive Range of Motion****Weeks 0 to 2:**

- Sling – for comfort only.

**Goals**

- Reduce inflammation.
- Decrease pain.
- Progression to full range of motion.
- Postural education.

**Manual therapy**

- Ice and modalities to reduce pain and inflammation.
- STM – global shoulder and CT junction.
- Graded GH mobilizations.
- ST mobilizations.

**Exercise progression**

- Progress to full range of motion, flexion and external rotation as tolerated.
  - Use a combination of wand, pulleys, wall walks or table slides to ensure compliance.
- Gradual introduction to internal rotation shoulder extensions (stick off back).
- Cervical range of motion and basic deep neck flexor activation (chin tucks).
- Active elbow, hand and wrist range of motion.
- Active shoulder retraction.
- Encourage walks and low intensity cardiovascular exercise to promote healing.

**Phase II – Progressive Stretching and Active Motion****Weeks 2 to 4-6:****Goals**

- Discontinue sling as instructed.
- Postural education.
- Full range of motion in all planes.

**Manual therapy**

- STM – global shoulder and CT junction.
- Scar tissue mobilization.
- Graded GH mobilizations.
- ST mobilizations.
- Gentle CR/RS for range of motion and RC-SS activation.

**Exercise progression**

- Progress to full range of motion using a combination of passive and active range of motion strategies.
- Serratus activation: ceiling punch (weight of arm) - may initially need assistance.
- Scapular strengthening – prone scapular series (rows and I's).
  - Emphasize scapular strengthening less than 90 degrees.
- External rotation on side (no resistance).
- Sub-maximal isometrics.
- Cervical range of motion as needed to maintain full mobility.
- DNF and proper postural positioning with all RC-SS exercises.
- Low to moderate cardiovascular work. May add elliptical but no running until 6 weeks.

---

**[www.meganwolfmd.com](http://www.meganwolfmd.com)**

**Phase III – Strengthening Phase****Weeks 4-6 to 12:**Goals

- Full active range of motion
- Normalize GH/ST arthrokinematics.
- Activate RC-SS with isometric and isotonic progression.

Manual therapy

- STM and joint mobilization to CT junction, GHJ and STJ as needed.
- CR/RS to gain range of motion while respecting repaired tissue.
- Manual perturbations.
- PNF patterns.

Exercise progression

- Continue with combined passive and active program to push full range of motion.
- Internal rotation with thumb up back and sleeper stretch.
- Continue with ceiling punch adding weight as tolerated.
- Sub-maximal rotator cuff isometrics (no pain).
- ER/IR isotonics at 0 degrees.
- Active ER at 90 degrees adding resistance as able.
- Advance prone series to include T's and Y's as tolerated.
- Add seated rows and front lat pulls.
- Biceps and triceps progressive resistance exercises (6-8 weeks BR and BT).
- Scaption: normalize ST arthrokinematics.
- CKC progression: quadruped, ball compression, counter weight shift, knee scapular push-ups, knee push-ups (all as tolerated).
  - Therapist directed RS and perturbations in quadruped – bilateral progressing to unilateral - tripod position.
- Supine progressing to standing PNF patterns, with resistance as appropriate.

**Phase IV – Advanced Strengthening and Plyometric Drills****Weeks 12 to 24:**Manual therapy

- STM and joint mobilization to CT junction, GHJ and STJ as needed.
- CR/RS to gain range of motion while respecting repaired tissue.
- Manual perturbations.
- PNF patterns.

Exercise progression (PRE/PSE)

- Full range of motion in all planes – emphasize terminal stretching.
- Advance strengthening at or above 90 degrees with prone or standing Y's and 90/90 as scapular control and ROM permit.
  - Patient health, physical condition and goals/objectives determine.
- Gym strengthening program: gradual progression with pressing and overhead activity.
- Progress closed kinetic chain program to include push-up progression beginning with counter, knee then gradual progression to full as appropriate.
- Continue with closed chain quadruped perturbations. Add open chain as strength permits.
- Initiate plyometric and rebounder drills as appropriate.

**Weeks 16 to 24:**Return to sport program

- Continue to progress RC and scapular strengthening program.

- Advance gym strengthening program.
- Return to sport testing for interval programs (golf, tennis etc.) using microfet dynamometer.
- Follow-up examination with the physician (4-6 months) for release to full activity.

Revised 9/2015